(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 13 December 2001 (13.12.2001)

PCT

(10) International Publication Number WO 01/93886 A1

(51) International Patent Classification7:

(21) International Application Number: I

PCT/EP00/05659

A61K 35/78

(22) International Filing Date:

19 June 2000 (19.06.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 00116395.7

8 June 2000 (08.06.2000) CN

(71) Applicants and

- (72) Inventors: ZHAO, Jian [CN/CN]: 420 Jiang Ning Road, 25E, Shanghai 200041 (CN). CHEN, Hu [CN/CN]; 420 Jiang Ning Road, 25E, Shanghai, 200041 (CN). JIN, Bei, Wen [CN/CN]; 420 Jiang Ning Road, 25E, Shanghai, 200041 (CN). ZHOU, Rui [CN/CN]; 420 Jiang Ning Road, 25E, Shanghai 200041 (CN).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): WOUTERS, Bert,
 C. [BE/BE]; Sysmans, Graaf, J. de Pret Staat 15, B-2900
 Schoten (BE).

- (74) Agent: SHOUSE, Emily, E.; Waddey & Patterson, 414 Union Street, Suite 2020, Nashville, TN 37219 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

A

(54) Title: METHOD OF OBTAINING A TEA PIGMENT FROM TEA LEAVES

(57) Abstract: The method according to the invention relates to a method of obtaining a tea pigment comprising theaflavin, thearubigin, theabrownin and catechins, characterised by (1) mixing tea leaves and ethanol, soaking and refluxing the obtained suspension; (2) centrifuging the suspension and discarding the pellet; (3) adding the remaining samples to a gel filtration column, washing the column; and (4) collecting the washing solution, extracting the tea pigment using a halogenated hydrocarbon having 1-3 carbon atoms, discarding the water phase, evaporating the halogenated hydrocarbon, and recovering a tea pigment powder containing less impurities and showing higher efficacy. The extraction product is applicable in the field of applications against hyperlipidemia and related diseases.

Method of obtaining a tea pigment from tea leaves

Description of the invention

The present invention provides a method of obtaining a tea pigment from tea

leaves. The obtained tea pigment comprises theaflavin, thearubigin, theabrownin and catechins.

Detailed description of the invention

The invention relates to a method comprising the following steps:

- 10 (1) mixing tea leaves and thanol, soaking and refluxing the obtained suspension;
 - (2) centrifuging the suspension and discarding the pellet;
 - (3) adding the remaining samples to a gel filtration column, washing the column; and
 - (4) collecting the washing solution, extracting the tea pigment using a halogenated hydrocarbon having 1 – 3 carbon atoms, discarding the water phase, evaporating the halogenated hydrocarbon and recovering a tea pigment powder containing less impurities and showing higher efficacy.

More in particular the method according to the invention comprises the following steps:

- (1) mixing tea leaves and ethanol in a w/w ratio of 1:1-20, preferably about 1:10, soaking and refluxing the obtained suspension;
- (2) centrifuging the suspension and discarding the pellet;
- (3) adding the samples to a Sephadex column, washing the column; and
- (4) collecting the washing solution, extracting tea pigment using a chlorinated hydrocarbon having one carbon atom, discarding the water phase, evaporating the chlorinated hydrocarbon and recovering a tea pigment powder containing less impurities and showing higher efficacy.

Most preferably the method according to the invention is carried out in the following way:

Mixing tea and 80% ethanol solution (w/w = 1:10), soaking the suspension at room temperature for 2 hours, heating and refluxing the suspension for 1 hour, filtrating the tea extraction solution, eliminating the tea residue; and

15

20

10

- 2. adjusting the suspension pH to 3.2, using 1mol/L HC1, 10,000 rpm centrifuge for 30 min, discarding the pellet; and
- adjusting the supernatant solution pH to 7.0, using 1mol/L NaOH, adding the sample to a Sephadex LH-20 column, washing the column with 40% - 100% ethanol solution; and
- 4. collecting the washing solution, adjusting the washing solution pH to 8.0, using 1mol/L NaOH, extracting tea pigment using CH₂Cl₂, discarding the water phase, evaporating the CH₂Cl₂ and obtaining a tea pigment powder having excellent pharmaceutical properties, in particular in applications against hyperlipidemia and related diseases.


Further the invention relates to the product obtained by means of the method defined above.

Another aspect of the convention relates to the pharmaceutical compositions comprising the product obtained by means of method defined above. The pharmaceutical composition is applicable in the field of applications against hyperlipidemia and related diseases.

Figure

20

The production process according to the invention is illustrated by the following figure.



CID: <WO_____0193886A1_I_>

10

4

Experimental Studies and Clinical Application

1. Animal experiments

(1) Safety test

Tea pigment obtained according to the method of the invention is poured directly into rats' stomachs. The dose for rats is 100 times higher than that for human. Observing these rats' behaviour for one week, we find that rats act normally. This test shows the tea pigment according to the invention is safe.

(2) Effect on serum lipids

Tea pigment according to the invention can induce decrease of serum total cholesterol (TC), triglyceride (TG), low density lipoprotein (LDL), and raise high density lipoprotein cholesterol (HDL) in comparison with control groups.




Figure 2. effects of tea pigment on serum lipids

15

2. Population Studies

(1) Hyperlipidemia study

The following values of TC, TG, HDL, LDL are the standard for normal human:

TC 3.1 - 5.7mol/L

TG 0.56 - 1.7mol/L

HDL 1.04 - 1.55mol/L

LDL 1.80 - 3.36mol/L

20

BNSDOCID: <WO____0193886A1_I_>

A total of 1,696 patients participated in this trial, 920 males and 776 females, ages 35 to 81 with an average weight of 58.9 ± 7.9 . Patients took 125 mg tea pigment 3 times/day for 4 weeks.

- 5 Patients satisfied the following criteria,
 - 1) did not suffer an acute heart attack, brain damage, no injury, no operations within the last 6 months;
 - 2) no kidney diseases;
 - 3) no diabetes mellitus;
- 10 4) no thyroid disease;
 - 5) no phase III hypertension;
 - 6) no drug induced hyperlipidemia;
 - 7) no pregnant women.
- 15 Table 1 changes in blood TC, TG, HDL and LDL levels.

	Cases (n)	Before treatment (mol/L)	After treatment (mol/L)	Rate of change (%)	P value
TC	811	6.71 + 0.55	5.65 + 0.41	- 15.8	< 0,01
TG	923	2.95 + 0.59	2.27 + 0.31	- 23.1	< 0,01
LDL	154	4.05 + 0.37	3.35 + 0.34	- 17.3	< 0,01
HDL	276	1.19 + 0.28	1.34 + 0.19	+ 12.6	< 0,01




Fig. 3 changes in patients blood TC, TG, HDL and LDL levels

20 Table 2 shows changes on patients with hyperlipidemia.

	Cases (n)	Markedly	Improved (%)	Not improved (%)	Total
İ	1	Improved (%)			improvement
TC	811	59.8 (485)	12.5 (101)	27.7 (225)	72.3
TG	923	41.7 (385)	24.1 (222)	34.2 (316)	65.8
LDL	154	48.7 (75)	31.2 (48)	20.1 (31)	79.9
HDL	276	50.7 (140)	23.9 (66)	25.4 (70)	74.7

(2) Dose-Effect and Period of Treatment-Effect Relationships:

250mg versus 125mg Tea pigment

60 days versus 30 days of Tea pigment treatment.

A total of 521 patients participated in this trial, 310 males and 211 females, ages 28 to 79 with an average weight of 55.2 ± 5.9 .

Group A: 125 mg 3 times a day

Group B: 250 mg 3 times a day

Patients satisfied the following criteria,

10

- did not suffer an acute heart attack, brain damage, no injury, no operations within last 6 months;
- 2) no kidney diseases;
- 3) no diabetes mellitus
- 4) no thyroid disease

15

20

- 5) no phase III hypertension
- 6) no drug induced hyperlipidemia
- 7) no pregnant women.

Table 3 shows changes in blood TC, TG, HDL and LDL levels. The period of treatment is 30 days.

ļ	Group	Cases (n)	Before treatment	After treatment	Rate of	P value
			(mol/L)	(mol/L)	change (%)	
TC	Α	210	6.51 ± 0.97	5.51 <u>+</u> 0.76	-15.3	<0,01
TC	В	156	6.54 ± 0.88	5.21 ± 0.61	-20.3	<0,01
TG	Α	198	3.11 ± 0.74	2.34 ± 0.46	-24.8	<0,01
TG	В	112	3.06 ± 0.78	2.01 ± 0.71	-34.3	<0,01
LDL	Α	165	4.32 <u>+</u> 0.76	3.41 ± 0.59	-21.1	<0,01
LDL	В	120	4.26 ± 0.81	3.03 ± 0.65	-28.9	<0,01
HDL	Α	171	1.04 ± 0.35	1.23 ± 0.29	+18.3	<0,01
HDL	В	104	0.98 ± 0.26	1.31 ± 0.22	+33.7	<0,01




Fig. 4 changes in patient's blood TC, TG, HDL and LDL levels (30 days)

Table 4 shows changes on patients with hyperlipidemia The period of treatment is 30 days.

	Group	Cases (n)	Markedly	Improved	Not Improved	Total
		_	Improved (%)	(%)	(%)	improvement %
TC	A	210	55.2 (116)	15.2 (32)	29.6 (62)	70.4
TC	В	156	62.8 (98)	14.1 (22)	23.1 (36)	76.9
TG	A	198	39.9 (79)	21.2 (42)	38.9 (77)	61.1
TG	В	112	50.9 (57)	19.6 (22)	29.5 (33)	70.5
LDL	A	165	47.3 (78)	28.5 (47)	24.2 (4)	75.8
LDL	В	120	50.8 (61)	30.0 (36)	19.2 (23)	81.8
HDL	A	171	49.7 (85)	21.6 (37)	28.7 (49)	71.4
HDL	В	104	55.8 (58)	24.0 (25)	20.2 (21)	79.8

Table 5 shows changes in blood TC, TG, HDL and LDL levels. The period of treatment is 60 days.

	Group	Cases (n)	Before treatment	After treatment	Rate of	P value
			(mol/L)	(mol/L)	change (%)	
TC	A	210	6.51 ± 0.97	5.48 ± 0.73	-15.8	<0,01
TC	В	156	6.54 ± 0.88	5.14 ± 0.75	-21.4	<0,01
TG	A	198	3.11 <u>+</u> 0.74	2.30 ± 0.46	-26.0	<0,01
TG	В	112	3.06 ± 0.78	1.92 ± 0.53	-37.2	<0,01
LDL	Α	165	4.32 <u>+</u> 0.76	3.38 ± 0.46	-21.8	<0,01
LDL	В	120	4.26 ± 0.81	2.89 ± 0.58	-32.2	<0,01
HDL	Α	171	1.04 ± 0.35	1.24 ± 0.24	+19.2	<0,01
HDL	В	104	0.98 ± 0.26	1.34 ± 0.19	+36.7	<0,01




Fig. 5 changes in atients' blood TC, TG, HDL and LDL levels (60 days).

Table 6 shows changes on patients with hyperlipidemia.

5 The period of treatment is 60 days.

	Group	Cases (n)	Markedly	Improved	Not Improved	Total
l	·		Improved (%)	(%)	(%)	improvement %
TC	Α	210	59.0 (124)	12.4 (26)	28.6 (60)	71.4
TC	В	156	67.9 (106)	10.3 (16)	21.8 (34)	78.2
TG	Α	198	46.5 (92)	17.6 (35)	35.9 (71)	64.1
TG	В	112	59.8 (67)	13.4 (15)	26.8 (30)	73.2
LDL	Α	165	57.0 (94)	20.0 (33)	23.0 (38)	77.0
LDL	В	120	65.0 (78)	18.3 (22)	16.7 (20)	83.3
HDL	Α	171	54.4 (93)	18.1 (31)	27.5 (47)	72.5
HDL	В	104	59.6 (62)	21.2 (22)	19.2 (20)	81.8

Significant differences occurred in all lipid level measurements.

However, no significant differences (P>0.05) occurred in the same dose and 2 periods of treatment.

10

Claims

10

15

- 1. Method of obtaining a tea pigment comprising theaflavin, thearubigin, theabrownin and catechins, characterised by
 - (1) mixing tea leaves and ethanol, soaking and refluxing the obtained suspension;
- 5 (2) centrifuging the suspension and discarding the pellet;
 - (3) adding the remaining samples to a gel filtration column, washing the column; and
 - (4) collecting the washing solution, extracting the tea pigment using a halogenated hydrocarbon having 1-3 carbon atoms, discarding the water phase, evaporating the halogenated hydrocarbon, and recovering a tea pigment powder containing less impurities and showing higher efficacy.
 - 2. Method according to claim 1, characterised in that in step (1) the tea leaves and the ethanol are mixed in a w/w-ratio of 1:1-20.
 - 3. Method according to claim 2, characterised in that in step (1) the tea leaves and the ethanol are mixed in a w/w-ratio of 1:10.
 - 4. Method according to any of claims 1-3, characterised in that in step (3) the samples are added to a Sephadex column, preferably a Sephadox LH-20 column.
 - 5. Method according to any of the claims 1-4, characterised in that in step (4) the halogenated hydrocarbon is a chlorinated hydrocarbon having one carbon atom.
- 20 6. Method according to claim 5 characterised in that in step (4) the halogenated hydrocarbon is CH₂Cl₂.
 - 7. The product obtained by carrying out the method according to any of the claims 1-
- 8. A pharmaceutical composition, comprising the product according to claim 7 as one of the active components.

DCID: <WO_____0193886A1_I_>

Fig 1




Fig 2




Fig 3




Fig 4





Fig 5



SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

national Application No

			101/EP 00/	705659
A. CLASSI IPC 7	IFICATION OF SUBJECT MATTER A61K35/78			
According to	o International Patent Classification (IPC) or to both national class	sification and IPC		
B. FIELDS	SEARCHED			
Minimum de IPC 7	ocumentation searched (classification system followed by classifi $A61K$	cation symbols)		
Documenta	lion searched other than minimum documentation to the extent th	at such documents are incl	uded in the fields sea	arched
Electronic d	fata base consulted during the international search (name of data	base and, where practical	l, search terms used)	
BIOSIS	, EPO-Internal, WPI Data, PAJ, FST	A, MEDLINE, PA	SCAL, CAB D	ata, EMBASE
C. DOCUM	ENTS CONSIDERED TO BE RELEVANT			
Category °	Citation of document, with indication, where appropriate, of the	relevant passages		Relevant to claim No.
X	US 4 613 672 A (HARA YUKIHIKO) 23 September 1986 (1986-09-23) column 2, line 20 -column 3, li	ne 59		1-8
A	DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE PHILADELPHIA, PA, US; 1992 HAYASHI M ET AL: "INHIBITORY EF GREEN TEA EXTRACTED WITH WATER ON SERUM LIPIDS OF RATS TREATED TRITON WR-1339" Database accession no. PREV1992 XP002164188 abstract	FECTS OF OR ETHANOL WITH		1,8
	& PHARMACOMETRICS, vol. 43, no. 6, 1992, pages 555 ISSN: 0300-8533 ———	5 -559 ,		
		-/		
X Furti	her documents are listed in the continuation of box C.	X Patent family	members are listed in	аплех.
"A" docume consider filing of the citation of the results of the citation of the results of the citation of the results of the	ent defining the general state of the art which is not dered to be of particular relevance document but published on or after the international tate ent which may throw doubts on priority ctaim(s) or is cited to establish the publication date of another n or other special reason (as specified) ent referring to an oral disclosure, use, exhibition or means ent published prior to the international filing date but han the priority date claimed	cited to understan invention "X" document of particle cannot be conside involve an invention "Y" document of particle cannot be considered document is combined to combine in the considered invention in the considered invention inventi	d not in conflict with the did the principle or the clared novel or cannot two step when the document to involve an involve an involve with one or morpination being obvious	the application but bory underlying the almed invention be considered to ument is taken atone almed invention entive step when the e other such docu- s to a person skilled
Date of the	actual completion of the international search	Date of mailing of	the international sear	ch report
1	0 July 2001	20/07/2	2001	_
Name and r	mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31–70) 340–2040, Tx. 31 651 epo nl,	Authorized officer		
	Fax: (+31-70) 340-3016	Rempp,	G	

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

national Application No

		rcT/EP 00/05659
	ntion) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	DATABASE WPI Section Ch, Week 198711 Derwent Publications Ltd., London, GB; Class B04, AN 1987-076886 XP002164189 & JP 62 030711 A (FURUKAWA NETSUGAKE), 9 February 1987 (1987-02-09) abstract	1,8
	·	
1		

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4613672	Α	23-09-1986	JP 1775152 C JP 2022755 B JP 60013780 A	28-07-1993 21-05-1990 24-01-1985
JP 62030711	Α	09-02-1987	NONE	

Form PCT/ISA/210 (patent family annex) (July 1992)